

PubMed Literature Search Categorized

Records from PubMed (www.pubmed.gov) – 2001-2005

Keywords used: Geographic Information Systems, GIS, Spatial Analysis

Literature search conducted 10/26/05 by Tara Murray, PRI Information Core

Literature search results categorized by Becki Curtis and Stephen Matthews

Health

Accidents

Grabowski, J. G., Curriero, F. C., Baker, S. P., & Li, G. (2002). Exploratory spatial analysis of pilot fatality rates in general aviation crashes using geographic information systems. *Am J Epidemiol*, 155(5), 398-405.

Grabowski, J. G., Curriero, F. C., Baker, S. P., & Li, G. (2002). Geographic patterns of pilot fatality rates in commuter and air taxi crashes. *Aviat Space Environ Med*, 73(10), 1014-1020.

Hijar, M., Trostle, J., & Bronfman, M. (2003). Pedestrian injuries in Mexico: a multi-method approach. *Soc Sci Med*, 57(11), 2149-2159.

Kam, B. H. (2003). A disaggregate approach to crash rate analysis. *Accid Anal Prev*, 35(5), 693-709.

Lightstone, A. S., Dhillon, P. K., Peek-Asa, C., & Kraus, J. F. (2001). A geographic analysis of motor vehicle collisions with child pedestrians in Long Beach, California: comparing intersection and midblock incident locations. *Inj Prev*, 7(2), 155-160.

Ng, K. S., Hung, W. T., & Wong, W. G. (2002). An algorithm for assessing the risk of traffic accident. *J Safety Res*, 33(3), 387-410.

Noland, R. B., & Quddus, M. A. (2004). A spatially disaggregate analysis of road casualties in England. *Accid Anal Prev*, 36(6), 973-984.

Rodriguez, D. Y., Fernandez, F. J., & Acero Velasquez, H. (2003). Road traffic injuries in Colombia. *Inj Control Saf Promot*, 10(1-2), 29-35.

Ruff, T. M., & Holden, T. P. (2003). Preventing collisions involving surface mining equipment: a GPS-based approach. *J Safety Res*, 34(2), 175-181.

Schulte Gary, S. L., Aultman-Hall, L., McCourt, M., & Stamatiadis, N. (2003). Consideration of driver home county prohibition and alcohol-related vehicle crashes. *Accid Anal Prev*, 35(5), 641-648.

Zhang, S. N., Lu, W., Zhou, S. F., Chen, S. M., Yang, B. J., Li, Q., et al. (2004). [Epidemiological study on traffic accident in Shanghai from 2000 to 2002]. *Zhonghua Liu Xing Bing Xue Za Zhi*, 25(3), 199-203.

Aging

Yiannakoulis, N., Rowe, B. H., Svenson, L. W., Schopflocher, D. P., Kelly, K., & Voaklander, D. C. (2003). Zones of prevention: the geography of fall injuries in the elderly. *Soc Sci Med*, 57(11), 2065-2073.

Alcohol

Gorman, D. M., Speer, P. W., Gruenewald, P. J., & Labouvie, E. W. (2001). Spatial dynamics of alcohol availability, neighborhood structure and violent crime. *J Stud Alcohol*, 62(5), 628-636.

Vanlaar, W. (2005). Drink driving in Belgium: results from the third and improved roadside survey. *Accid Anal Prev*, 37(3), 391-397.

Weitzman, E. R., Folkman, A., Folkman, M. P., & Wechsler, H. (2003). The relationship of alcohol outlet density to heavy and frequent drinking and drinking-related problems among college students at eight universities. *Health Place*, 9(1), 1-6.

Child Health

Andrade, A. L., Silva, S. A., Martelli, C. M., Oliveira, R. M., Morais Neto, O. L., Siqueira Junior, J. B., et al. (2004). Population-based surveillance of pediatric pneumonia: use of spatial analysis in an urban area of Central Brazil. *Cad Saude Publica*, 20(2), 411-421.

Brooker, S., Beasley, M., Ndinaromtan, M., Madjiouroum, E. M., Baboguel, M., Djenguinabe, E., et al. (2002). Use of remote sensing and a geographical information system in a national helminth control programme in Chad. *Bull World Health Organ*, 80(10), 783-789.

Chakraborty, J. (2001). Analyzing exposure of school children to accidental releases of hazardous substances. *J Expo Anal Environ Epidemiol*, 11(4), 269-278.

Chen, Y., Waters, E., & Green, J. (2002). Geospatial analysis of childhood pertussis in Victoria, 1993-97. *Aust N Z J Public Health*, 26(5), 456-461.

Gething, P. W., Noor, A. M., Zurovac, D., Atkinson, P. M., Hay, S. I., Nixon, M. S., et al. (2004). Empirical modelling of government health service use by children with fevers in Kenya. *Acta Trop*, 91(3), 227-237.

Hruba, F., Fabianova, E., Koppova, K., & Vandenberg, J. J. (2001). Childhood respiratory symptoms, hospital admissions, and long-term exposure to airborne particulate matter. *J Expo Anal Environ Epidemiol*, *11*(1), 33-40.

Lercher, P., Evans, G. W., Meis, M., & Kofler, W. W. (2002). Ambient neighbourhood noise and children's mental health. *Occup Environ Med*, *59*(6), 380-386.

Loyola, E., Castillo-Salgado, C., Najera-Aguilar, P., Vidaurre, M., Mujica, O. J., & Martinez-Piedra, R. (2002). [Geographic information systems as a tool for monitoring health inequalities]. *Rev Panam Salud Publica*, *12*(6), 415-428.

Neto, O. L., Barros, M. B., Martelli, C. M., Silva, S. A., Cavenaghi, S. M., & Siqueira, J. B., Jr. (2001). [Differential patterns of neonatal and post-neonatal mortality rates in Goiania, Brazil, 1992-1996: use of spatial analysis to identify high-risk areas]. *Cad Saude Publica*, *17*(5), 1241-1250.

Birth Defects

Wu, J., Wang, J., Meng, B., Chen, G., Pang, L., Song, X., et al. (2004). Exploratory spatial data analysis for the identification of risk factors to birth defects. *BMC Public Health*, *4*(1), 23.

Yauck, J. S., Malloy, M. E., Blair, K., Simpson, P. M., & McCarver, D. G. (2004). Proximity of residence to trichloroethylene-emitting sites and increased risk of offspring congenital heart defects among older women. *Birth Defects Res A Clin Mol Teratol*, *70*(10), 808-814.

Diet/Obesity

Block, J. P., Scribner, R. A., & DeSalvo, K. B. (2004). Fast food, race/ethnicity, and income: a geographic analysis. *Am J Prev Med*, *27*(3), 211-217.

Brown, D. (2004). Mapping the future of dietetics with geographic information systems. *J Am Diet Assoc*, *104*(5), 727-728.

Burdette, H. L., & Whitaker, R. C. (2004). Neighborhood playgrounds, fast food restaurants, and crime: relationships to overweight in low-income preschool children. *Prev Med*, *38*(1), 57-63.

Liu, G. C., Cunningham, C., Downs, S. M., Marrero, D. G., & Fineberg, N. (2002). A spatial analysis of obesogenic environments for children. *Proc AMIA Symp*, 459-463.

Mobley, L. R., Finkelstein, E. A., Khavjou, O. A., & Will, J. C. (2004). Spatial analysis of body mass index and smoking behavior among WISEWOMAN participants. *J Womens Health (Larchmt)*, *13*(5), 519-528.

Zenk, S. N., Schulz, A. J., Israel, B. A., James, S. A., Bao, S., & Wilson, M. L. (2005). Neighborhood racial composition, neighborhood poverty, and the spatial accessibility of supermarkets in metropolitan Detroit. *Am J Public Health, 95*(4), 660-667.

Disease

Ali, M., Emch, M., Yunus, M., & Sack, R. B. (2001). Are the environmental niches of *Vibrio cholerae* O139 different from those of *Vibrio cholerae* O1 El Tor? *Int J Infect Dis, 5*(4), 214-219.

Bakker, M. I., Hatta, M., Kwenang, A., Faber, W. R., van Beers, S. M., Klatser, P. R., et al. (2004). Population survey to determine risk factors for *Mycobacterium leprae* transmission and infection. *Int J Epidemiol, 33*(6), 1329-1336.

Barcellos, C., & Sabroza, P. C. (2001). The place behind the case: leptospirosis risks and associated environmental conditions in a flood-related outbreak in Rio de Janeiro. *Cad Saude Publica, 17 Suppl*, 59-67.

Baumgardner, D. J., Steber, D., Glazier, R., Paretsky, D. P., Egan, G., Baumgardner, A. M., et al. (2005). Geographic information system analysis of blastomycosis in northern Wisconsin, USA: waterways and soil. *Med Mycol, 43*(2), 117-125.

Bavia, M. E., Carneiro, D. D., Gurgel Hda, C., Madureira Filho, C., & Barbosa, M. G. (2005). Remote Sensing and Geographic Information Systems and risk of American visceral leishmaniasis in Bahia, Brazil. *Parassitologia, 47*(1), 165-169.

Bergquist, N. R. (2001). Vector-borne parasitic diseases: new trends in data collection and risk assessment. *Acta Trop, 79*(1), 13-20.

Bernardi, M. (2001). Linkages between FAO agroclimatic data resources and the development of GIS models for control of vector-borne diseases. *Acta Trop, 79*(1), 21-34.

Boccia, D., Oliver, C. I., Charlett, A., Bennett, S., Orr, H., Sarangi, J., et al. (2004). Outbreak of a new *Salmonella* phage type in South West England: alternative epidemiological investigations are needed. *Commun Dis Public Health, 7*(4), 339-343.

Brianti, E., Malone, J. B., McCarroll, J. C., Bernardi, M., Drigo, M., & Gruszynski, K. (2004). [A minimum medical GIS database (MMDb) for Europe]. *Parassitologia, 46*(1-2), 67-70.

Brooker, S., Kabatereine, N. B., Tukahebwa, E. M., & Kazibwe, F. (2004). Spatial analysis of the distribution of intestinal nematode infections in Uganda. *Epidemiol Infect, 132*(6), 1065-1071.

Brownstein, J. S., Rosen, H., Purdy, D., Miller, J. R., Merlino, M., Mostashari, F., et al. (2002). Spatial analysis of West Nile virus: rapid risk assessment of an introduced vector-borne zoonosis. *Vector Borne Zoonotic Dis, 2*(3), 157-164.

- Bunnell, J. E., Price, S. D., Das, A., Shields, T. M., & Glass, G. E. (2003). Geographic information systems and spatial analysis of adult *Ixodes scapularis* (Acari: Ixodidae) in the Middle Atlantic region of the U.S.A. *J Med Entomol*, 40(4), 570-576.
- Cattand, P., Jannin, J., & Lucas, P. (2001). Sleeping sickness surveillance: an essential step towards elimination. *Trop Med Int Health*, 6(5), 348-361.
- Chaput, E. K., Meek, J. I., & Heimer, R. (2002). Spatial analysis of human granulocytic ehrlichiosis near Lyme, Connecticut. *Emerg Infect Dis*, 8(9), 943-948.
- Cifuentes, E., Mazari-Hiriart, M., Carneiro, F., Bianchi, F., & Gonzalez, D. (2002). The risk of enteric diseases in young children and environmental indicators in sentinel areas of Mexico City. *Int J Environ Health Res*, 12(1), 53-62.
- Cortinas, M. R., Guerra, M. A., Jones, C. J., & Kitron, U. (2002). Detection, characterization, and prediction of tick-borne disease foci. *Int J Med Microbiol*, 291 Suppl 33, 11-20.
- Cromley, E. K. (2003). GIS and disease. *Annu Rev Public Health*, 24, 7-24.
- Danson, F. M., Graham, A. J., Pleydell, D. R., Campos-Ponce, M., Giraudoux, P., & Craig, P. S. (2003). Multi-scale spatial analysis of human alveolar echinococcosis risk in China. *Parasitology*, 127 Suppl, S133-141.
- Elnaiem, D. E., Schorscher, J., Bendall, A., Obsomer, V., Osman, M. E., Mekkawi, A. M., et al. (2003). Risk mapping of visceral leishmaniasis: the role of local variation in rainfall and altitude on the presence and incidence of kala-azar in eastern Sudan. *Am J Trop Med Hyg*, 68(1), 10-17.
- Fang, L. Q., Cao, W. C., Chen, H. X., Wang, B. G., Wu, X. M., Yang, H., et al. (2003). [Study on the application of geographic information system in spatial distribution of hemorrhage fever with renal syndrome in China]. *Zhonghua Liu Xing Bing Xue Za Zhi*, 24(4), 265-268.
- Fang, L. Q., Cao, W. C., Dun, Z., Wu, X. M., Sun, P. Y., Kulldorff, M., et al. (2003). [Spatial analysis on the distribution of hemorrhagic fever with renal syndrome by geographic information system in Haidian district, Beijing]. *Zhonghua Liu Xing Bing Xue Za Zhi*, 24(11), 1020-1023.
- Forand, S. P., Talbot, T. O., Druschel, C., & Cross, P. K. (2002). Data quality and the spatial analysis of disease rates: congenital malformations in New York State. *Health Place*, 8(3), 191-199.
- Fuentes, M. V., Malone, J. B., & Mas-Coma, S. (2001). Validation of a mapping and prediction model for human fasciolosis transmission in Andean very high altitude endemic areas using remote sensing data. *Acta Trop*, 79(1), 87-95.
- Fuentes, M. V., Sainz-Elipse, S., Nieto, P., Malone, J. B., & Mas-Coma, S. (2005). Geographical Information Systems risk assessment models for zoonotic fascioliasis in the South American Andes region. *Parassitologia*, 47(1), 151-156.

- Gatton, M. L., Kelly-Hope, L. A., Kay, B. H., & Ryan, P. A. (2004). Spatial-temporal analysis of Ross River virus disease patterns in Queensland, Australia. *Am J Trop Med Hyg*, 71(5), 629-635.
- Gebre-Michael, T., Malone, J. B., & McNally, K. (2005). Use of Geographic Information Systems in the development of prediction models for onchocerciasis control in Ethiopia. *Parassitologia*, 47(1), 135-144.
- Glavanakov, S., White, D. J., Caraco, T., Lapenis, A., Robinson, G. R., Szymanski, B. K., et al. (2001). Lyme disease in New York State: spatial pattern at a regional scale. *Am J Trop Med Hyg*, 65(5), 538-545.
- Guerra, M., Walker, E., Jones, C., Paskewitz, S., Cortinas, M. R., Stancil, A., et al. (2002). Predicting the risk of Lyme disease: habitat suitability for *Ixodes scapularis* in the north central United States. *Emerg Infect Dis*, 8(3), 289-297.
- Hassan, A. N. (2004). Bancroftian filariasis: spatial patterns, environmental correlates and landscape predictors of disease risk. *J Egypt Soc Parasitol*, 34(2), 501-513.
- Henrich, T. J., Hutchaleelaha, S., Jiwariyavej, V., Barbazan, P., Nitatpattana, N., Yoksan, S., et al. (2003). Geographic dynamics of viral encephalitis in Thailand. *Microbes Infect*, 5(7), 603-611.
- Jarup, L. (2004). Health and environment information systems for exposure and disease mapping, and risk assessment. *Environ Health Perspect*, 112(9), 995-997.
- Jon, Z. (2001). Methods of mortality (incidence) rates interpretation assessment in small populations. *Cent Eur J Public Health*, 9(1), 14-21.
- Krech, T. (2002). TBE foci in Switzerland. *Int J Med Microbiol*, 291 Suppl 33, 30-33.
- Lai, P. C., Wong, C. M., Hedley, A. J., Lo, S. V., Leung, P. Y., Kong, J., et al. (2004). Understanding the spatial clustering of severe acute respiratory syndrome (SARS) in Hong Kong. *Environ Health Perspect*, 112(15), 1550-1556.
- Le Bras, M., & Malvy, J. M. (2004). [The tropical pathogenic complex: a new look at an old concept]. *Med Trop (Mars)*, 64(6), 613-618.
- Lewis, M. D., Pavlin, J. A., Mansfield, J. L., O'Brien, S., Boomsma, L. G., Elbert, Y., et al. (2002). Disease outbreak detection system using syndromic data in the greater Washington DC area. *Am J Prev Med*, 23(3), 180-186.
- Lointier, M., Truc, P., Drapeau, L., Nanga, S., & Tarek, M. (2001). [Methodology to determine risk zones for sleeping sickness in Cote d'Ivoire by the spatial approach]. *Med Trop (Mars)*, 61(4-5), 390-396.

- Malone, J. B. (2005). Biology-based mapping of vector-borne parasites by Geographic Information Systems and Remote Sensing. *Parassitologia*, 47(1), 27-50.
- Malone, J. B., McNally, K. L., McCarroll, J. C., Corbett, J. D., & Mkoji, G. (2004). Modeling the biocoenose of parasitic diseases using remote sensing and geographic information systems. *Parassitologia*, 46(1-2), 59-61.
- Montenegro, A. C., Werneck, G. L., Kerr-Pontes, L. R., Barreto, M. L., & Feldmeier, H. (2004). Spatial analysis of the distribution of leprosy in the State of Ceara, Northeast Brazil. *Mem Inst Oswaldo Cruz*, 99(7), 683-686.
- Morrison, A. C., Gray, K., Getis, A., Astete, H., Sihuincha, M., Focks, D., et al. (2004). Temporal and geographic patterns of *Aedes aegypti* (Diptera: Culicidae) production in Iquitos, Peru. *J Med Entomol*, 41(6), 1123-1142.
- Munch, Z., Van Lill, S. W., Booyesen, C. N., Zietsman, H. L., Enarson, D. A., & Beyers, N. (2003). Tuberculosis transmission patterns in a high-incidence area: a spatial analysis. *Int J Tuberc Lung Dis*, 7(3), 271-277.
- Noma, M., Nwoke, B. E., Nutall, I., Tambala, P. A., Enyong, P., Namsenmo, A., et al. (2002). Rapid epidemiological mapping of onchocerciasis (REMO): its application by the African Programme for Onchocerciasis Control (APOC). *Ann Trop Med Parasitol*, 96 Suppl 1, S29-39.
- Nyamogoba, H. D., Obala, A. A., & Kakai, R. (2002). Combating cholera epidemics by targeting reservoirs of infection and transmission routes: a review. *East Afr Med J*, 79(3), 150-155.
- Nygard, K., Andersson, Y., Rottingen, J. A., Svensson, A., Lindback, J., Kistemann, T., et al. (2004). Association between environmental risk factors and campylobacter infections in Sweden. *Epidemiol Infect*, 132(2), 317-325.
- Odiit, M., Coleman, P. G., McDermott, J. J., Fevre, E. M., Welburn, S. C., & Woolhouse, M. E. (2004). Spatial and temporal risk factors for the early detection of *Trypanosoma brucei* rhodesiense sleeping sickness patients in Tororo and Busia districts, Uganda. *Trans R Soc Trop Med Hyg*, 98(10), 569-576.
- Park, B. J., Sigel, K., Vaz, V., Komatsu, K., McRill, C., Phelan, M., et al. (2005). An epidemic of coccidioidomycosis in Arizona associated with climatic changes, 1998-2001. *J Infect Dis*, 191(11), 1981-1987.
- Polack, S. R., Solomon, A. W., Alexander, N. D., Massae, P. A., Safari, S., Shao, J. F., et al. (2005). The household distribution of trachoma in a Tanzanian village: an application of GIS to the study of trachoma. *Trans R Soc Trop Med Hyg*, 99(3), 218-225.
- Pugliatti, M., Solinas, G., Sotgiu, S., Castiglia, P., & Rosati, G. (2002). Multiple sclerosis distribution in northern Sardinia: spatial cluster analysis of prevalence. *Neurology*, 58(2), 277-282.

Reynolds, P., Elkin, E., Scalf, R., Von Behren, J., & Neutra, R. R. (2001). A case-control pilot study of traffic exposures and early childhood leukemia using a geographic information system. *Bioelectromagnetics, Suppl 5*, S58-68.

Richardson, M., van Lill, S. W., van der Spuy, G. D., Munch, Z., Booysen, C. N., Beyers, N., et al. (2002). Historic and recent events contribute to the disease dynamics of Beijing-like *Mycobacterium tuberculosis* isolates in a high incidence region. *Int J Tuberc Lung Dis*, 6(11), 1001-1011.

Rissland, J., van Treeck, U., Taeger, D., & Baumeister, H. G. (2003). [Infection surveillance in North Rhine-Westphalia--standard reports, barometer and early warning system]. *Gesundheitswesen*, 65(12), 719-723.

Rodriguez, A. P., Serpa, I. M., & Garcia, M. (2002). [Meningococcal disease in the City of Havana before and after vaccination]. *Rev Cubana Med Trop*, 54(2), 106-112.

Rogers, D. J., & Randolph, S. E. (2003). Studying the global distribution of infectious diseases using GIS and RS. *Nat Rev Microbiol*, 1(3), 231-237.

Rytönen, M., Ranta, J., Tuomilehto, J., & Karvonen, M. (2001). Bayesian analysis of geographical variation in the incidence of Type I diabetes in Finland. *Diabetologia*, 44 Suppl 3, B37-44.

Saathoff, E., Olsen, A., Kvalsvig, J. D., Appleton, C. C., Sharp, B., & Kleinschmidt, I. (2005). Ecological covariates of *Ascaris lumbricoides* infection in schoolchildren from rural KwaZulu-Natal, South Africa. *Trop Med Int Health*, 10(5), 412-422.

Saathoff, E., Olsen, A., Sharp, B., Kvalsvig, J. D., Appleton, C. C., & Kleinschmidt, I. (2005). Ecologic covariates of hookworm infection and reinfection in rural Kwazulu-natal/south Africa: a geographic information system-based study. *Am J Trop Med Hyg*, 72(4), 384-391.

Samuelsson, U., & Lofman, O. (2004). Geographical mapping of type 1 diabetes in children and adolescents in south east Sweden. *J Epidemiol Community Health*, 58(5), 388-392.

Schlundt, D. G., Mushi, C., Larson, C. O., & Marrs, M. (2001). Use of innovative technologies in the evaluation of Nashville's REACH 2010 community action plan: reducing disparities in cardiovascular disease and diabetes in the African American community. *J Ambul Care Manage*, 24(3), 51-60.

Seketeli, A., Adeoye, G., Eyamba, A., Nnoruka, E., Drameh, P., Amazigo, U. V., et al. (2002). The achievements and challenges of the African Programme for Onchocerciasis Control (APOC). *Ann Trop Med Parasitol*, 96 Suppl 1, S15-28.

Serpa, I. M., Rodriguez, A. P., & Garcia, M. (2001). [Bacterial meningitis: geographic and referential study in the City of Havana in 1998]. *Rev Cubana Med Trop*, 53(3), 204-211.

- Simoes, L. B., Marques, S. A., & Bagagli, E. (2004). Distribution of paracoccidioidomycosis: determination of ecologic correlates through spatial analyses. *Med Mycol*, 42(6), 517-523.
- Stein, K. J., & Claborn, D. M. (2005). Telephonic survey of surveillance and control procedures for the mosquito vectors of West Nile virus near naval installations in the eastern United States. *Mil Med*, 170(8), 658-662.
- Tong, S., Bi, P., Hayes, J., Donald, K., & Mackenzie, J. (2001). Geographic variation of notified Ross River virus infections in Queensland, Australia, 1985-1996. *Am J Trop Med Hyg*, 65(3), 171-176.
- Tong, S., Hayes, J. F., & Dale, P. (2005). Spatiotemporal variation of notified Barmah Forest virus infections in Queensland, Australia, 1993-2001. *Int J Environ Health Res*, 15(2), 89-98.
- Toubiana, L., Richard, J. B., & Landais, P. (2005). Geographical information system for end-stage renal disease: SIGNe, an aid to public health decision making. *Nephrol Dial Transplant*, 20(2), 273-277.
- Uphoff, H., Stalleicken, I., Bartelds, A., Phiesel, B., & Kistemann, B. T. (2004). Are influenza surveillance data useful for mapping presentations? *Virus Res*, 103(1-2), 35-46.
- Vieira, V., Webster, T., Aschengrau, A., & Ozonoff, D. (2002). A method for spatial analysis of risk in a population-based case-control study. *Int J Hyg Environ Health*, 205(1-2), 115-120.
- Viik-Kajander, M., Moltchanova, E., Salomaa, V., Tuomilehto, J., Ketonen, M., Palomaki, P., et al. (2003). Geographical variation in the incidence of acute myocardial infarction in eastern Finland--a Bayesian perspective. *Ann Med*, 35(1), 43-50.
- Werneck, G. L., Costa, C. H., Walker, A. M., David, J. R., Wand, M., & Maguire, J. H. (2002). The urban spread of visceral leishmaniasis: clues from spatial analysis. *Epidemiology*, 13(3), 364-367.
- Williams, C. J., Willocks, L. J., Lake, I. R., & Hunter, P. R. (2004). Geographic correlation between deprivation and risk of meningococcal disease: an ecological study. *BMC Public Health*, 4(1), 30.
- Yabsley, M. J., Wimberly, M. C., Stallknecht, D. E., Little, S. E., & Davidson, W. R. (2005). Spatial analysis of the distribution of Ehrlichia chaffeensis, causative agent of human monocytotropic ehrlichiosis, across a multi-state region. *Am J Trop Med Hyg*, 72(6), 840-850.
- Yelifari, L., Bloch, P., Magnussen, P., van Lieshout, L., Dery, G., Anemana, S., et al. (2005). Distribution of human Oesophagostomum bifurcum, hookworm and Strongyloides stercoralis infections in northern Ghana. *Trans R Soc Trop Med Hyg*, 99(1), 32-38.

Zeman, P., & Bene, C. (2004). A tick-borne encephalitis ceiling in Central Europe has moved upwards during the last 30 years: possible impact of global warming? *Int J Med Microbiol*, 293 Suppl 37, 48-54.

Zenilman, J. M., Glass, G., Shields, T., Jenkins, P. R., Gaydos, J. C., & McKee, K. T., Jr. (2002). Geographic epidemiology of gonorrhoea and chlamydia on a large military installation: application of a GIS system. *Sex Transm Infect*, 78(1), 40-44.

Asthma

Allacci, M. S. (2005). Identifying environmental risk factors for asthma emergency care" a multilevel approach for ecological study. *J Ambul Care Manage*, 28(1), 2-15.

Lewis, S. A., Antoniak, M., Venn, A. J., Davies, L., Goodwin, A., Salfield, N., et al. (2005). Secondhand smoke, dietary fruit intake, road traffic exposures, and the prevalence of asthma: a cross-sectional study in young children. *Am J Epidemiol*, 161(5), 406-411.

Oyana, T. J., Rogerson, P., & Lwebuga-Mukasa, J. S. (2004). Geographic clustering of adult asthma hospitalization and residential exposure to pollution at a United States-Canada border crossing. *Am J Public Health*, 94(7), 1250-1257.

Vegni, F. E., Panceri, M. L., & Wilkinson, P. (2003). [Antiasthmatic drug consumption as an indicator of the prevalence of respiratory pathology in a pediatric population]. *Ann Ig*, 15(4), 311-317.

Cancer

Aldrich, T. E., Andrews, K. W., & Liboff, A. R. (2001). Brain cancer risk and electromagnetic fields (EMFs): assessing the geomagnetic component. *Arch Environ Health*, 56(4), 314-319.

Bell, B. S. (2002). Spatial analysis of disease--applications. *Cancer Treat Res*, 113, 151-182.

Brody, J. G., Aschengrau, A., McKelvey, W., Rudel, R. A., Swartz, C. H., & Kennedy, T. (2004). Breast cancer risk and historical exposure to pesticides from wide-area applications assessed with GIS. *Environ Health Perspect*, 112(8), 889-897.

Chelule, P. K., Gqaleni, N., Dutton, M. F., & Chuturgoon, A. A. (2001). Exposure of rural and urban populations in KwaZulu Natal, South Africa, to fumonisin B(1) in maize. *Environ Health Perspect*, 109(3), 253-256.

- Fano, V., Michelozzi, P., Ancona, C., Capon, A., Forastiere, F., & Perucci, C. A. (2004). Occupational and environmental exposures and lung cancer in an industrialised area in Italy. *Occup Environ Med*, 61(9), 757-763.
- Floret, N., Mauny, F., Challier, B., Cahn, J. Y., Tourneux, F., & Viel, J. F. (2004). [Dioxin emissions and soft-tissue sarcoma: results of a population-based case-control study]. *Rev Epidemiol Sante Publique*, 52(3), 213-220.
- Floret, N., Mauny, F., Challier, B., Arveux, P., Cahn, J. Y., & Viel, J. F. (2003). Dioxin emissions from a solid waste incinerator and risk of non-Hodgkin lymphoma. *Epidemiology*, 14(4), 392-398.
- French, J. L., & Wand, M. P. (2004). Generalized additive models for cancer mapping with incomplete covariates. *Biostatistics*, 5(2), 177-191.
- Menegozzo, M., Trinca, S., Cammino, F., Mastrantonio, M., Menegozzo, S., Sturchio, A., et al. (2004). [Geographical distribution of mortality from malignant pleural neoplasms and of former asbestos-exposed workers in the Campania Region]. *Epidemiol Prev*, 28(3), 150-155.
- O'Leary, E. S., Vena, J. E., Freudenheim, J. L., & Brasure, J. (2004). Pesticide exposure and risk of breast cancer: a nested case-control study of residentially stable women living on Long Island. *Environ Res*, 94(2), 134-144.
- Paldy, A., Pinter, A., Nador, G., Vincze, I., & Malnasi, T. (2003). [Regional differences of mortality from malignancies in Hungary]. *Orv Hetil*, 144(25), 1227-1233.
- Poulstrup, A., & Hansen, H. L. (2004). Use of GIS and exposure modeling as tools in a study of cancer incidence in a population exposed to airborne dioxin. *Environ Health Perspect*, 112(9), 1032-1036.
- Reynolds, P., Hurley, S. E., Goldberg, D. E., Yerabati, S., Gunier, R. B., Hertz, A., et al. (2004). Residential proximity to agricultural pesticide use and incidence of breast cancer in the California Teachers Study cohort. *Environ Res*, 96(2), 206-218.
- Roche, L. M., Skinner, R., & Weinstein, R. B. (2002). Use of a geographic information system to identify and characterize areas with high proportions of distant stage breast cancer. *J Public Health Manag Pract*, 8(2), 26-32.
- Sandor, J., Szerencse, P., Szucs, M., Nemeth, A., Kiss, I., & Ember, I. (2003). [Investigation of regional clusters of environmentally induced cancers]. *Magy Onkol*, 47(2), 177-183.
- Scott, D., Curtis, B., & Twumasi, F. O. (2002). Towards the creation of a health information system for cancer in KwaZulu-Natal, South Africa. *Health Place*, 8(4), 237-249.

Selvin, S., & Merrill, D. W. (2002). Adult leukemia: a spatial analysis. *Epidemiology*, 13(2), 151-156.

Speer, S. A., Semenza, J. C., Kurosaki, T., & Anton-Culver, H. (2002). Risk factors for acute myeloid leukemia and multiple myeloma: a combination of GIS and case-control studies. *J Environ Health*, 64(7), 9-16; quiz 35-16.

Tempest, V., Higgs, G., McDonald, K., Iredale, R., Bater, T., & Gray, J. (2005). A pilot study of spatial patterns in referrals to a multicentre cancer genetics service. *Community Genet*, 8(2), 73-79.

Verkasalo, P. K., Kokki, E., Pukkala, E., Vartiainen, T., Kiviranta, H., Penttinen, A., et al. (2004). Cancer risk near a polluted river in Finland. *Environ Health Perspect*, 112(9), 1026-1031.

Wang, F. (2004). Spatial clusters of cancers in Illinois 1986-2000. *J Med Syst*, 28(3), 237-256.

Dengue Fever

Ali, M., Wagatsuma, Y., Emch, M., & Breiman, R. F. (2003). Use of a geographic information system for defining spatial risk for dengue transmission in Bangladesh: role for *Aedes albopictus* in an urban outbreak. *Am J Trop Med Hyg*, 69(6), 634-640.

Barcellos, C., Pustai, A. K., Weber, M. A., & Brito, M. R. (2005). [Identification of places with potential transmission of dengue fever in Porto Alegre using Geographical Information Systems]. *Rev Soc Bras Med Trop*, 38(3), 246-250.

Chadee, D. D., Williams, F. L., & Kitron, U. D. (2005). Impact of vector control on a dengue fever outbreak in Trinidad, West Indies, in 1998. *Trop Med Int Health*, 10(8), 748-754.

Tran, A., DeParis, X., Dussart, P., Morvan, J., Rabarison, P., Remy, F., et al. (2004). Dengue spatial and temporal patterns, French Guiana, 2001. *Emerg Infect Dis*, 10(4), 615-621.

Van Benthem, B. H., Vanwambeke, S. O., Khantikul, N., Burghoorn-Maas, C., Panart, K., Oskam, L., et al. (2005). Spatial patterns of and risk factors for seropositivity for dengue infection. *Am J Trop Med Hyg*, 72(2), 201-208.

HIV/AIDS

Desjeux, P., & Alvar, J. (2003). Leishmania/HIV co-infections: epidemiology in Europe. *Ann Trop Med Parasitol*, 97 Suppl 1, 3-15.

Desjeux, P., Piot, B., O'Neill, K., & Meert, J. P. (2001). [Co-infections of leishmania/HIV in south Europe]. *Med Trop (Mars)*, 61(2), 187-193.

Fulcher, C., & Kaukinen, C. (2005). Mapping and visualizing the location HIV service providers: an exploratory spatial analysis of Toronto neighborhoods. *AIDS Care*, 17(3), 386-396.

Fulcher, C. L., & Kaukinen, C. E. (2003). HIV/AIDS Community Health Information System. *AMIA Annu Symp Proc*, 1064.

Torugsa, K., Anderson, S., Thongsen, N., Sirisopana, N., Jugsudee, A., Junlananto, P., et al. (2003). HIV epidemic among young Thai Men, 1991-2000. *Emerg Infect Dis*, 9(7), 881-883.

Welton, A., Adelberger, K., Patterson, K., & Gilbert, D. (2004). Optimal placement of syringe-exchange programs. *J Urban Health*, 81(2), 268-277.

Malaria

Brooker, S., Clarke, S., Njagi, J. K., Polack, S., Mugo, B., Estambale, B., et al. (2004). Spatial clustering of malaria and associated risk factors during an epidemic in a highland area of western Kenya. *Trop Med Int Health*, 9(7), 757-766.

Ceccato, P., Connor, S. J., Jeanne, I., & Thomson, M. C. (2005). Application of Geographical Information Systems and Remote Sensing technologies for assessing and monitoring malaria risk. *Parassitologia*, 47(1), 81-96.

Claborn, D. M., Masuoka, P. M., Klein, T. A., Hooper, T., Lee, A., & Andre, R. G. (2002). A cost comparison of two malaria control methods in Kyunggi Province, Republic of Korea, using remote sensing and geographic information systems. *Am J Trop Med Hyg*, 66(6), 680-685.

Gemperli, A., Vounatsou, P., Kleinschmidt, I., Bagayoko, M., Lengeler, C., & Smith, T. (2004). Spatial patterns of infant mortality in Mali: the effect of malaria endemicity. *Am J Epidemiol*, 159(1), 64-72.

Hassan, A. N., Kenawy, M. A., Kamal, H., Abdel Sattar, A. A., & Sowilem, M. M. (2003). GIS-based prediction of malaria risk in Egypt. *East Mediterr Health J*, 9(4), 548-558.

Hay, S. I., Guerra, C. A., Tatem, A. J., Noor, A. M., & Snow, R. W. (2004). The global distribution and population at risk of malaria: past, present, and future. *Lancet Infect Dis*, 4(6), 327-336.

- Keating, J., MacIntyre, K., Mbogo, C., Githeko, A., Regens, J. L., Swalm, C., et al. (2003). A geographic sampling strategy for studying relationships between human activity and malaria vectors in urban Africa. *Am J Trop Med Hyg*, 68(3), 357-365.
- Kleinschmidt, I., Sharp, B. L., Clarke, G. P., Curtis, B., & Fraser, C. (2001). Use of generalized linear mixed models in the spatial analysis of small-area malaria incidence rates in Kwazulu Natal, South Africa. *Am J Epidemiol*, 153(12), 1213-1221.
- Kleinschmidt, I., Sharp, B., Mueller, I., & Vounatsou, P. (2002). Rise in malaria incidence rates in South Africa: a small-area spatial analysis of variation in time trends. *Am J Epidemiol*, 155(3), 257-264.
- Leonardo, L. R., Rivera, P. T., Crisostomo, B. A., Sarol, J. N., Bantayan, N. C., Tiu, W. U., et al. (2005). A study of the environmental determinants of malaria and schistosomiasis in the Philippines using Remote Sensing and Geographic Information Systems. *Parassitologia*, 47(1), 105-114.
- Martin, C., Curtis, B., Fraser, C., & Sharp, B. (2002). The use of a GIS-based malaria information system for malaria research and control in South Africa. *Health Place*, 8(4), 227-236.
- Mnzava, A. E., Sharp, B. L., Mthembu, D. J., le Sueur, D., Dlamini, S. S., Gumede, J. K., et al. (2001). Malaria control--two years' use of insecticide-treated bednets compared with insecticide house spraying in KwaZulu-Natal. *S Afr Med J*, 91(11), 978-983.
- Morrison, A. C., Astete, H., Chapilliquen, F., Ramirez-Prada, C., Diaz, G., Getis, A., et al. (2004). Evaluation of a sampling methodology for rapid assessment of *Aedes aegypti* infestation levels in Iquitos, Peru. *J Med Entomol*, 41(3), 502-510.
- Najera-Aguilar, P., Martinez-Piedra, R., & Vidaurre-Arenas, M. (2005). From sketch to digital maps: a Geographic Information System (GIS) Model and Application for malaria control without the use of pesticides. *Epidemiol Bull*, 26(1), 11.
- Nihei, N., Hashida, Y., Kobayashi, M., & Ishii, A. (2002). Analysis of malaria endemic areas on the Indochina Peninsula using remote sensing. *Jpn J Infect Dis*, 55(5), 160-166.
- Noor, A. M., Zurovac, D., Hay, S. I., Ochola, S. A., & Snow, R. W. (2003). Defining equity in physical access to clinical services using geographical information systems as part of malaria planning and monitoring in Kenya. *Trop Med Int Health*, 8(10), 917-926.
- Phillips-Howard, P. A., ter Kuile, F. O., Nahlen, B. L., Alaii, J. A., Gimnig, J. E., Kolczak, M. S., et al. (2003). The efficacy of permethrin-treated bed nets on child mortality and morbidity in western Kenya II. Study design and methods. *Am J Trop Med Hyg*, 68(4 Suppl), 10-15.

Rakotomanana, F., Jeanne, I., Duchemin, J. B., Pietra, V., Raharimalala, L., Tombo, M. L., et al. (2001). [Geographic approach in malaria control in the central highlands of Madagascar]. *Arch Inst Pasteur Madagascar*, 67(1-2), 27-30.

Singer, B. H., & de Castro, M. C. (2001). Agricultural colonization and malaria on the Amazon frontier. *Ann N Y Acad Sci*, 954, 184-222.

Sithiprasasna, R., Linthicum, K. J., Liu, G. J., Jones, J. W., & Singhasivanon, P. (2003). Some entomological observations on temporal and spatial distribution of malaria vectors in three villages in northwestern Thailand using a geographic information system. *Southeast Asian J Trop Med Public Health*, 34(3), 505-516.

Sithiprasasna, R., Ugsang, D. M., Honda, K., Jones, J. W., & Singhasivanon, P. (2005). Ikonos-derived malaria transmission risk in northwestern Thailand. *Southeast Asian J Trop Med Public Health*, 36(1), 14-22.

Socheat, D., Denis, M. B., Fandeur, T., Zhang, Z., Yang, H., Xu, J., et al. (2003). Mekong malaria. II. Update of malaria, multi-drug resistance and economic development in the Mekong region of Southeast Asia. *Southeast Asian J Trop Med Public Health*, 34 Suppl 4, 1-102.

Srivastava, A., Nagpal, B. N., Saxena, R., Eapen, A., Ravindran, K. J., Subbarao, S. K., et al. (2003). GIS based malaria information management system for urban malaria scheme in India. *Comput Methods Programs Biomed*, 71(1), 63-75.

Su, Y. Q., Zhang, Z. Y., Xu, D. Z., Xi, Y. Z., Wang, S. Q., & Li, C. X. (2003). [Study on the distribution of malaria in Hainan province through spatial local interpolation technique]. *Zhonghua Liu Xing Bing Xue Za Zhi*, 24(4), 269-271.

Yang, G. J., Zhou, X. N., Malone, J. B., McCarroll, J. C., Wang, T. P., & Liu, J. X. (2002). [Application of multifactor spatial composite model to predict transmission tendency of malaria at national level]. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi*, 20(3), 145-147.

Yang, G., Zhou, X., Malone, J. B., McCarroll, J. C., Wang, T., Liu, J., et al. (2002). GIS prediction model of malaria transmission in Jiangsu province. *Zhonghua Yu Fang Yi Xue Za Zhi*, 36(2), 103-105.

Yu, G. W., & Tang, L. H. (2004). [Remote sensing and geographic information system techniques in malaria research]. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi*, 22(2), 119-121.

Schistosomiasis

- Abdel-Rahman, M. S., El-Bahy, M. M., Malone, J. B., Thompson, R. A., & El Bahy, N. M. (2001). Geographic information systems as a tool for control program management for schistosomiasis in Egypt. *Acta Trop*, 79(1), 49-57.
- Barbosa, C. S., Araujo, K. C., Antunes, L., Favre, T., & Pieri, O. S. (2004). Spatial distribution of schistosomiasis foci on Itamaraca Island, Pernambuco, Brazil. *Mem Inst Oswaldo Cruz*, 99(5 Suppl 1), 79-83.
- Bavia, M. E., Malone, J. B., Hale, L., Dantas, A., Marroni, L., & Reis, R. (2001). Use of thermal and vegetation index data from earth observing satellites to evaluate the risk of schistosomiasis in Bahia, Brazil. *Acta Trop*, 79(1), 79-85.
- Brooker, S. (2002). Schistosomes, snails and satellites. *Acta Trop*, 82(2), 207-214.
- Clennon, J. A., King, C. H., Muchiri, E. M., Kariuki, H. C., Ouma, J. H., Mungai, P., et al. (2004). Spatial patterns of urinary schistosomiasis infection in a highly endemic area of coastal Kenya. *Am J Trop Med Hyg*, 70(4), 443-448.
- Davis, G. M., Wu, W. P., Chen, H. G., Liu, H. Y., Guo, J. G., Lin, D. D., et al. (2002). A baseline study of importance of bovines for human *Schistosoma japonicum* infections around Poyang Lake, China: villages studied and snail sampling strategy. *Am J Trop Med Hyg*, 66(4), 359-371.
- Handzel, T., Karanja, D. M., Addiss, D. G., Hightower, A. W., Rosen, D. H., Colley, D. G., et al. (2003). Geographic distribution of schistosomiasis and soil-transmitted helminths in Western Kenya: implications for anthelmintic mass treatment. *Am J Trop Med Hyg*, 69(3), 318-323.
- Malone, J. B., Bergquist, N. R., Huh, O. K., Bavia, M. E., Bernardi, M., El Bahy, M. M., et al. (2001). A global network for the control of snail-borne disease using satellite surveillance and geographic information systems. *Acta Trop*, 79(1), 7-12.
- Malone, J. B., Yilma, J. M., McCarroll, J. C., Erko, B., Mukaratirwa, S., & Zhou, X. (2001). Satellite climatology and the environmental risk of *Schistosoma mansoni* in Ethiopia and east Africa. *Acta Trop*, 79(1), 59-72.
- Nihei, N., Kajihara, N., Kirinoki, M., Chigusa, Y., Saitoh, Y., Shimamura, R., et al. (2004). Fixed-point observation of *Oncomelania nosophora* in Kofu Basin -- establishment of monitoring system of schistosomiasis japonica in Japan. *Parasitol Int*, 53(2), 199-205.
- Raso, G., Matthys, B., N'Goran, E. K., Tanner, M., Vounatsou, P., & Utzinger, J. (2005). Spatial risk prediction and mapping of *Schistosoma mansoni* infections among schoolchildren living in western Cote d'Ivoire. *Parasitology*, 131(Pt 1), 97-108.

Stensgaard, A., Jorgensen, A., Kabatereine, N. B., Malone, J. B., & Kristensen, T. K. (2005). Modeling the distribution of *Schistosoma mansoni* and host snails in Uganda using satellite sensor data and Geographical Information Systems. *Parassitologia*, 47(1), 115-125.

Yang, G. J., Vounatsou, P., Zhou, X. N., Tanner, M., & Utzinger, J. (2005). A Bayesian-based approach for spatio-temporal modeling of county level prevalence of *Schistosoma japonicum* infection in Jiangsu province, China. *Int J Parasitol*, 35(2), 155-162.

Zhou, X. N., Malone, J. B., Kristensen, T. K., & Bergquist, N. R. (2001). Application of geographic information systems and remote sensing to schistosomiasis control in China. *Acta Trop*, 79(1), 97-106.

Drug Use

Fujiwara, T., Takano, T., & Nakamura, K. (2005). The spread of drug abuse in rapidly urbanizing communities in Vientiane, Lao People's Democratic Republic. *Health Promot Int*, 20(1), 61-68.

Mason, M., Cheung, I., & Walker, L. (2004). Substance use, social networks, and the geography of urban adolescents. *Subst Use Misuse*, 39(10-12), 1751-1777.

Emergency Management

Gall, M. (2004). Where to go? Strategic modelling of access to emergency shelters in Mozambique. *Disasters*, 28(1), 82-97.

Guidry, V. T., & Margolis, L. H. (2005). Unequal respiratory health risk: using GIS to explore hurricane-related flooding of schools in Eastern North Carolina. *Environ Res*, 98(3), 383-389.

Gutierrez, E., Taucer, F., De Groeve, T., Al-Khudhairy, D. H., & Zaldivar, J. M. (2005). Analysis of worldwide earthquake mortality using multivariate demographic and seismic data. *Am J Epidemiol*, 161(12), 1151-1158.

Kaiser, R., Spiegel, P. B., Henderson, A. K., & Gerber, M. L. (2003). The application of geographic information systems and global positioning systems in humanitarian emergencies: lessons learned, programme implications and future research. *Disasters*, 27(2), 127-140.

Keramitsoglou, I., Kiranoudis, C. T., Sarimveis, H., & Sifakis, N. (2004). A multidisciplinary decision support system for forest fire crisis management. *Environ Manage*, 33(2), 212-225.

Lindsey, J. (2003). GPS, computers, ambulances & getting lost. What do they have in common? *Jems*, 28(1), 91.

Mathew, D. (2005). Information technology and public health management of disasters--a model for South Asian countries. *Prehospital Disaster Med*, 20(1), 54-60.

Ott, W. E. (2003). Electronic & data management issues: global positioning system & EMS. *Jems*, 28(8), 94-97.

Peleg, K., & Pliskin, J. S. (2004). A geographic information system simulation model of EMS: reducing ambulance response time. *Am J Emerg Med*, 22(3), 164-170.

Verter, V., & Kara, B. Y. (2001). A GIS-based framework for hazardous materials transport risk assessment. *Risk Anal*, 21(6), 1109-1120.

Wang, X. Y., Qu, J. Y., Shi, Z. Q., & Ling, Y. S. (2003). A GIS-based prediction and assessment system of off-site accident consequence for Guangdong nuclear power plant. *Radiat Prot Dosimetry*, 106(3), 257-262.

Waring, S., Zakos-Feliberti, A., Wood, R., Stone, M., Padgett, P., & Arafat, R. (2005). The utility of geographic information systems (GIS) in rapid epidemiological assessments following weather-related disasters: methodological issues based on the Tropical Storm Allison Experience. *Int J Hyg Environ Health*, 208(1-2), 109-116.

Waring, S. C., Reynolds, K. M., D'Souza, G., & Arafat, R. R. (2002). Rapid assessment of household needs in the Houston area after Tropical Storm Allison. *Disaster Manag Response*, 3-9.

Williams, K. G., Schootman, M., Quayle, K. S., Struthers, J., & Jaffe, D. M. (2003). Geographic variation of pediatric burn injuries in a metropolitan area. *Acad Emerg Med*, 10(7), 743-752.

Epidemiology

Blaasaas, K. G., & Tynes, T. (2002). Comparison of three different ways of measuring distances between residences and high voltage power lines. *Bioelectromagnetics*, 23(4), 288-291.

Botto, C., Escalona, E., Vivas-Martinez, S., Behm, V., Delgado, L., & Coronel, P. (2005). Geographical patterns of onchocerciasis in southern Venezuela: relationships between environment and infection prevalence. *Parassitologia*, 47(1), 145-150.

Cringoli, G., Ippolito, A., & Taddei, R. (2005). Advances in satellite remote sensing of phenoclimatic features for epidemiological applications. *Parassitologia*, 47(1), 51-62.

Elliott, P., & Wartenberg, D. (2004). Spatial epidemiology: current approaches and future challenges. *Environ Health Perspect*, 112(9), 998-1006.

Hellstrom, L., Jarup, L., Persson, B., & Axelson, O. (2004). Using environmental concentrations of cadmium and lead to assess human exposure and dose. *J Expo Anal Environ Epidemiol*, *14*(5), 416-423.

Hoque, E., Hope, V., Scragg, R., Baker, M., & Shrestha, R. (2004). A descriptive epidemiology of giardiasis in New Zealand and gaps in surveillance data. *N Z Med J*, *117*(1205), U1149.

Idowu, E. T., Adewale, B., Mafe, M. A., Appelt, B., & Bamgbose, A. (2004). Endemicity of onchocerciasis in some local government areas of Niger state. *Afr J Med Med Sci*, *33*(1), 31-34.

Kabatereine, N. B., Brooker, S., Tukahebwa, E. M., Kazibwe, F., & Onapa, A. W. (2004). Epidemiology and geography of *Schistosoma mansoni* in Uganda: implications for planning control. *Trop Med Int Health*, *9*(3), 372-380.

Karsenty, E., & Leventhal, A. (2002). [Health Geographic Information System (HGIS)--a tool for health planning and epidemiology]. *Harefuah*, *141*(12), 1070-1075, 1089.

Kistemann, T., Zimmer, S., Vagsholm, I., & Andersson, Y. (2004). GIS-supported investigation of human EHEC and cattle VTEC O157 infections in Sweden: geographical distribution, spatial variation and possible risk factors. *Epidemiol Infect*, *132*(3), 495-505.

Krieger, N. (2003). Place, space, and health: GIS and epidemiology. *Epidemiology*, *14*(4), 384-385.

Martinez, R., Vidaurre, M., Najera, P., Loyola, E., Castillo-Salgado, C., & Eisner, C. (2001). SIGEpi: geographic information system in epidemiology and public health. *Epidemiol Bull*, *22*(3), 4-5.

Nuckols, J. R., Ward, M. H., & Jarup, L. (2004). Using geographic information systems for exposure assessment in environmental epidemiology studies. *Environ Health Perspect*, *112*(9), 1007-1015.

Rinaldi, L., Cascone, C., Sibilio, G., Musella, V., Taddei, R., & Cringoli, G. (2004). [Geographical Information Systems and remote sensing technologies in parasitological epidemiology]. *Parassitologia*, *46*(1-2), 71-74.

Health

Abellana, R. M., & Ascaso, C. (2004). [Statistical methods to analyze risk with spatial distribution patterns]. *Med Clin (Barc)*, *122* Suppl 1, 68-72.

Ali, M., Emch, M., Ashley, C., & Streatfield, P. K. (2001). Implementation of a medical geographic information system: concepts and uses. *J Health Popul Nutr*, *19*(2), 100-110.

- Bauer, I. L., & Puotinen, M. (2002). Geographic Information Systems and travel health. *J Travel Med*, 9(6), 308-314.
- Blanco, G. A., & Cooper, E. L. (2004). Immune systems, geographic information systems (GIS), environment and health impacts. *J Toxicol Environ Health B Crit Rev*, 7(6), 465-480.
- Caley, L. M. (2004). Using geographic information systems to design population-based interventions. *Public Health Nurs*, 21(6), 547-554.
- Castillo-Salgado, C., & Loyola, E. (2002). Development of the Healthy Condition Index using geographic information systems in health. *Epidemiol Bull*, 23(4), 7-11.
- Cockings, S., & Martin, D. (2005). Zone design for environment and health studies using pre-aggregated data. *Soc Sci Med*, 60(12), 2729-2742.
- Fu, H., Umezaki, M., Nakamura, K., Chen, P., Chen, J., Shi, H. J., et al. (2004). Pilot study of using GIS to visualize health status distribution: case study of Songjiang District, Shanghai. *J Med Dent Sci*, 51(4), 179-185.
- Fulcher, C. L., & Kaukinen, C. E. (2004). Visualizing the infrastructure of US healthcare using Internet GIS: a community health informatics approach for reducing health disparities. *Medinfo*, 11(Pt 2), 1197-1201.
- Harcourt, S. E., Edwards, D. E., Fleming, D. M., Smith, R. L., & Smith, G. E. (2004). How representative is the population covered by the RCGP spotter practice scheme? Using Geographical Information Systems to assess. *J Public Health (Oxf)*, 26(1), 88-94.
- Herbreteau, V., Salem, G., Souris, M., Hugot, J. P., & Gonzalez, J. P. (2005). Sizing up human health through remote sensing: uses and misuses. *Parassitologia*, 47(1), 63-79.
- Higgs, G., Smith, D. P., & Gould, M. I. (2005). Findings from a survey on GIS use in the UK National Health Service: organisational challenges and opportunities. *Health Policy*, 72(1), 105-117.
- Jenks, R. H., & Malecki, J. M. (2004). GIS--a proven tool for public health analysis. *J Environ Health*, 67(3), 32-34.
- Lerner, E. B., Fairbanks, R. J., & Shah, M. N. (2005). Identification of out-of-hospital cardiac arrest clusters using a geographic information system. *Acad Emerg Med*, 12(1), 81-84.
- McLafferty, S. L. (2003). GIS and health care. *Annu Rev Public Health*, 24, 25-42.
- Olvingson, C., Hallberg, J., Timpka, T., & Lindqvist, K. (2002). Ethical issues in public health informatics: implications for system design when sharing geographic information. *J Biomed Inform*, 35(3), 178-185.

Ricketts, T. C. (2003). Geographic information systems and public health. *Annu Rev Public Health, 24*, 1-6.

Health Care Service

Brabyn, L., & Barnett, R. (2004). Population need and geographical access to general practitioners in rural New Zealand. *N Z Med J, 117*(1199), U996.

Costa, L. S., Nassi, C. D., Pinheiro, R. S., & Almeida, R. M. (2003). Accessibility of selected hospitals and medical procedures by means of aerial and transit network-based measures. *Health Serv Manage Res, 16*(2), 136-140.

Faruque, F., Hewlett, P. O., Wyatt, S., Wilson, K., Lofton, S., Frate, D., et al. (2004). Geospatial information technology: an adjunct to service-based outreach and education. *J Nurs Educ, 43*(2), 88-91.

Foley, R. (2002). Assessing the applicability of GIS in a health and social care setting: planning services for informal carers in East Sussex, England. *Soc Sci Med, 55*(1), 79-96.

Gallegos, G. (2002). Estimating populations for collective dose calculations. *Health Phys, 83*(2), 283-286.

Ghinea, G., Gill, D., Frank, A., & de Souza, L. H. (2002). Using geographical information systems for management of back-pain data. *J Manag Med, 16*(2-3), 219-237.

Glazier, R. H., Creatore, M. I., Gozdyra, P., Matheson, F. I., Steele, L. S., Boyle, E., et al. (2004). Geographic methods for understanding and responding to disparities in mammography use in Toronto, Canada. *J Gen Intern Med, 19*(9), 952-961.

Heard, N. J., Larsen, U., & Hozumi, D. (2004). Investigating access to reproductive health services using GIS: proximity to services and the use of modern contraceptives in Malawi. *Afr J Reprod Health, 8*(2), 164-179.

Higgs, G., & Gould, M. (2001). Is there a role for GIS in the 'new NHS'? *Health Place, 7*(3), 247-259.

Hokerberg, Y. H., Duchiate, M. P., & Barcellos, C. (2001). [Organization and quality of health care for Kaingang Indians in Rio Grande do Sul, Brazil]. *Cad Saude Publica, 17*(2), 261-272.

Juarez, P. D., Robinson, P. L., & Matthews-Juarez, P. (2002). 100% access, zero health disparities, and GIS: an improved methodology for designating health professions shortage areas. *J Health Soc Policy, 16*(1-2), 155-167.

Kaneko, Y., Takano, T., & Nakamura, K. (2003). Visual localisation of community health needs to rational decision-making in public health services. *Health Place, 9*(3), 241-251.

- Kitajima, T., Kitazawa, T., Ing, C. K., & Noyama, O. (2001). [An attempt to estimate geographical accessibility to day service facilities among the elderly with a geographical information system]. *Nippon Koshu Eisei Zasshi*, 48(8), 613-619.
- Kumar, N. (2004). Changing geographic access to and locational efficiency of health services in two Indian districts between 1981 and 1996. *Soc Sci Med*, 58(10), 2045-2067.
- Lin, S. J. (2004). Access to community pharmacies by the elderly in Illinois: a geographic information systems analysis. *J Med Syst*, 28(3), 301-309.
- Lovett, A., Haynes, R., Sunnenberg, G., & Gale, S. (2002). Car travel time and accessibility by bus to general practitioner services: a study using patient registers and GIS. *Soc Sci Med*, 55(1), 97-111.
- Luo, W. (2004). Using a GIS-based floating catchment method to assess areas with shortage of physicians. *Health Place*, 10(1), 1-11.
- Luo, W., Wang, F., & Douglass, C. (2004). Temporal changes of access to primary health care in Illinois (1990-2000) and policy implications. *J Med Syst*, 28(3), 287-299.
- McGregor, J., Hanlon, N., Emmons, S., Voaklander, D., & Kelly, K. (2005). If all ambulances could fly: putting provincial standards of emergency care access to the test in Northern British Columbia. *Can J Rural Med*, 10(3), 163-168.
- McLafferty, S., & Grady, S. (2004). Prenatal care need and access: a GIS analysis. *J Med Syst*, 28(3), 321-333.
- McLafferty, S., & Grady, S. (2005). Immigration and geographic access to prenatal clinics in Brooklyn, NY: a geographic information systems analysis. *Am J Public Health*, 95(4), 638-640.
- Moss, M. P., & Schell, M. C. (2004). GIS(c): a scientific framework and methodological tool for nursing research. *ANS Adv Nurs Sci*, 27(2), 150-159.
- Murad, A. A. (2004). Creating a GIS application for local health care planning in Saudi Arabia. *Int J Environ Health Res*, 14(3), 185-199.
- Noor, A. M., Gikandi, P. W., Hay, S. I., Muga, R. O., & Snow, R. W. (2004). Creating spatially defined databases for equitable health service planning in low-income countries: the example of Kenya. *Acta Trop*, 91(3), 239-251.
- O'Neill, L. (2003). Estimating out-of-hospital mortality due to myocardial infarction. *Health Care Manag Sci*, 6(3), 147-154.
- Parchman, M. L., Ferrer, R. L., & Blanchard, K. S. (2002). Geography and geographic information systems in family medicine research. *Fam Med*, 34(2), 132-137.

- Plescia, M., Koontz, S., & Laurent, S. (2001). Community assessment in a vertically integrated health care system. *Am J Public Health, 91*(5), 811-814.
- Rafalski, E., & Zun, L. (2004). Using GIS to monitor emergency room use in a large urban hospital in Chicago. *J Med Syst, 28*(3), 311-319.
- Randremanana, R. V., Migliani, R., Rakotomanga, S., & Jeanne, I. (2001). [Geographic Information Systems and health: application to Antananarivo city]. *Arch Inst Pasteur Madagascar, 67*(1-2), 74-78.
- Riner, M. E., Cunningham, C., & Johnson, A. (2004). Public health education and practice using geographic information system technology. *Public Health Nurs, 21*(1), 57-65.
- Rosero-Bixby, L. (2004). Spatial access to health care in Costa Rica and its equity: a GIS-based study. *Soc Sci Med, 58*(7), 1271-1284.
- Tanser, F. C. (2002). The application of GIS technology to equitably distribute fieldworker workload in a large, rural South African health survey. *Trop Med Int Health, 7*(1), 80-90.
- Tanser, F., Hosegood, V., Benzler, J., & Solarsh, G. (2001). New approaches to spatially analyse primary health care usage patterns in rural South Africa. *Trop Med Int Health, 6*(10), 826-838.
- Wang, F., & Luo, W. (2005). Assessing spatial and nonspatial factors for healthcare access: towards an integrated approach to defining health professional shortage areas. *Health Place, 11*(2), 131-146.
- Wood, D. J., Clark, D., & Gatrell, A. C. (2004). Equity of access to adult hospice inpatient care within north-west England. *Palliat Med, 18*(6), 543-549.
- Yu, W., Cowper, D., Berger, M., Kuebeler, M., Kubal, J., & Manheim, L. (2004). Using GIS to profile health-care costs of VA Quality-Enhancement Research Initiative diseases. *J Med Syst, 28*(3), 271-285.

Dentistry

- Bartling, W. C., & Schleyer, T. K. (2003). An application of Geospatial Information System (GIS) technology to anatomic dental charting. *AMIA Annu Symp Proc, 786*.
- Higgs, G., & Richards, W. (2002). The use of geographical information systems in examining variations in sociodemographic profiles of dental practice catchments: a case study of a Swansea practice. *Prim Dent Care, 9*(2), 63-69.
- Krause, D., Frate, D. A., & May, W. L. (2005). Demographics and distribution of dentists in Mississippi: a dental work force study. *J Am Dent Assoc, 136*(5), 668-677.

Marriott, A., Harding, W., Devlin, N., & Benwell, G. (2001). The delivery of orthodontic care in New Zealand. Part I: Analysis of a census of orthodontists. *N Z Dent J*, 97(429), 87-92.

Susi, L., & Mascarenhas, A. K. (2002). Using a geographical information system to map the distribution of dentists in Ohio. *J Am Dent Assoc*, 133(5), 636-642.

Mental Health

Goldenberg, J. S., & Contreras Escudero, L. (2002). [Design and implementation of an epidemiological surveillance system for mental health]. *Rev Panam Salud Publica*, 11(2), 83-92.

Mortality

Altavista, P., Belli, S., Bianchi, F., Binazzi, A., Comba, P., Del Giudice, R., et al. (2004). [Cause-specific mortality in an area of Campania with numerous waste disposal sites]. *Epidemiol Prev*, 28(6), 311-321.

Ferrandiz, J., Abellan, J. J., Gomez-Rubio, V., Lopez-Quilez, A., Sanmartin, P., Abellan, C., et al. (2004). Spatial analysis of the relationship between mortality from cardiovascular and cerebrovascular disease and drinking water hardness. *Environ Health Perspect*, 112(9), 1037-1044.

Mitchell, R., Dorling, D., & Shaw, M. (2002). Population production and modelling mortality--an application of geographic information systems in health inequalities research. *Health Place*, 8(1), 15-24.

Rytkonen, M., Rusanen, J., & Nayha, S. (2001). Small-area variation in mortality in the city of Oulu, Finland, during the period 1978--1995. *Health Place*, 7(2), 75-79.

Sankoh, O. A., Ye, Y., Sauerborn, R., Muller, O., & Becher, H. (2001). Clustering of childhood mortality in rural Burkina Faso. *Int J Epidemiol*, 30(3), 485-492.

Physical Activity

Duncan, M., & Mummery, K. (2005). Psychosocial and environmental factors associated with physical activity among city dwellers in regional Queensland. *Prev Med*, 40(4), 363-372.

Estabrooks, P. A., Lee, R. E., & Gyurcsik, N. C. (2003). Resources for physical activity participation: does availability and accessibility differ by neighborhood socioeconomic status? *Ann Behav Med*, 25(2), 100-104.

- Frank, L. D., Schmid, T. L., Sallis, J. F., Chapman, J., & Saelens, B. E. (2005). Linking objectively measured physical activity with objectively measured urban form: findings from SMARTRAQ. *Am J Prev Med*, 28(2 Suppl 2), 117-125.
- King, W. C., Belle, S. H., Brach, J. S., Simkin-Silverman, L. R., Soska, T., & Kriska, A. M. (2005). Objective measures of neighborhood environment and physical activity in older women. *Am J Prev Med*, 28(5), 461-469.
- Kirtland, K. A., Porter, D. E., Addy, C. L., Neet, M. J., Williams, J. E., Sharpe, P. A., et al. (2003). Environmental measures of physical activity supports: perception versus reality. *Am J Prev Med*, 24(4), 323-331.
- Leslie, E., Saelens, B., Frank, L., Owen, N., Bauman, A., Coffee, N., et al. (2005). Residents' perceptions of walkability attributes in objectively different neighbourhoods: a pilot study. *Health Place*, 11(3), 227-236.
- Li, F., Fisher, K. J., Brownson, R. C., & Bosworth, M. (2005). Multilevel modelling of built environment characteristics related to neighbourhood walking activity in older adults. *J Epidemiol Community Health*, 59(7), 558-564.
- Makikawa, M., Isaka, T., Iida, T., Asajima, S., Tsukise, K., Shiozawa, N., et al. (2004). Jogging support system with portable monitoring device and health manage software. *Medinfo*, 11(Pt 2), 1260-1263.
- Pikora, T. J., Bull, F. C., Jamrozik, K., Knuiaman, M., Giles-Corti, B., & Donovan, R. J. (2002). Developing a reliable audit instrument to measure the physical environment for physical activity. *Am J Prev Med*, 23(3), 187-194.
- Troped, P. J., Saunders, R. P., Pate, R. R., Reininger, B., Ureda, J. R., & Thompson, S. J. (2001). Associations between self-reported and objective physical environmental factors and use of a community rail-trail. *Prev Med*, 32(2), 191-200.
- Wilson, D. K., Kirtland, K. A., Ainsworth, B. E., & Addy, C. L. (2004). Socioeconomic status and perceptions of access and safety for physical activity. *Ann Behav Med*, 28(1), 20-28.

Environment

Agriculture

Colby, M. M., & Johnson, Y. J. (2002). Potential uses for geographic information system-based planning and decision support technology in intensive food animal production. *Anim Health Res Rev*, 3(1), 31-42.

Dalgaard, T., Kjeldsen, C., Hutchings, N. J., & Hansen, J. F. (2001). N-losses and energy use in a scenario for conversion to organic farming. *ScientificWorldJournal*, 1 Suppl 2, 822-829.

Vela Acosta, M. S., Reding, D. J., Cooper, S. P., & Gunderson, P. (2005). Lessons learned: geographic information systems and farmworkers in the Lake States. *J Agric Saf Health*, 11(1), 85-97.

Wing, S., Freedman, S., & Band, L. (2002). The potential impact of flooding on confined animal feeding operations in eastern North Carolina. *Environ Health Perspect*, 110(4), 387-391.

Animal Studies

Barrera, R., Torres, N., Freier, J. E., Navarro, J. C., Garcia, C. Z., Salas, R., et al. (2001). Characterization of enzootic foci of Venezuelan equine encephalitis virus in western Venezuela. *Vector Borne Zoonotic Dis*, 1(3), 219-230.

Environment and Land Use

Angelstam, P., Boresjo-Bronge, L., Mikusinski, G., Sporrang, U., & Wastfelt, A. (2003). Assessing village authenticity with satellite images: a method to identify intact cultural landscapes in Europe. *Ambio*, 32(8), 594-604.

Brown, M. T., & Vivas, M. B. (2005). Landscape development intensity index. *Environ Monit Assess*, 101(1-3), 289-309.

Danz, N. P., Regal, R. R., Niemi, G. J., Brady, V. J., Hollenhorst, T., Johnson, L. B., et al. (2005). Environmentally stratified sampling design for the development of Great Lakes environmental indicators. *Environ Monit Assess*, 102(1-3), 41-65.

Guan, Y. N., Guo, S., Xue, Y., Li, R., & Yang, Q. K. (2003). Landscape dynamic change in Mu Us Desert derived from Landsat TM data. *J Environ Sci (China)*, 15(2), 222-226.

Hassan, M. M., Atkins, P. J., & Dunn, C. E. (2003). The spatial pattern of risk from arsenic poisoning: a Bangladesh case study. *J Environ Sci Health A Tox Hazard Subst Environ Eng*, 38(1), 1-24.

- Heitgerd, J. L., & Lee, V. C. (2003). A new look at neighborhoods near National Priorities List sites. *Soc Sci Med*, 57(6), 1117-1126.
- Hellmeier, W., Queste, A., & Woltering, R. (2001). [Surveillance of drinking water supply with a geographic information system: a pilot project of the of the Iogd NRM and the Hoxter district]. *Gesundheitswesen*, 63(3), 170-175.
- Jerrett, M., Burnett, R. T., Goldberg, M. S., Sears, M., Krewski, D., Catalan, R., et al. (2003). Spatial analysis for environmental health research: concepts, methods, and examples. *J Toxicol Environ Health A*, 66(16-19), 1783-1810.
- Kistemann, T., Dangendorf, F., & Schweikart, J. (2002). New perspectives on the use of Geographical Information Systems (GIS) in environmental health sciences. *Int J Hyg Environ Health*, 205(3), 169-181.
- Kitsiou, D., Coccossis, H., & Karydis, M. (2002). Multi-dimensional evaluation and ranking of coastal areas using GIS and multiple criteria choice methods. *Sci Total Environ*, 284(1-3), 1-17.
- Krause, G., Bock, M., Weiers, S., & Braun, G. (2004). Mapping land-cover and mangrove structures with remote sensing techniques: a contribution to a synoptic GIS in support of coastal management in North Brazil. *Environ Manage*, 34(3), 429-440.
- Kristensen, T. K., Malone, J. B., & McCarroll, J. C. (2001). Use of satellite remote sensing and geographic information systems to model the distribution and abundance of snail intermediate hosts in Africa: a preliminary model for *Biomphalaria pfeifferi* in Ethiopia. *Acta Trop*, 79(1), 73-78.
- Leimgruber, P., Christen, C. A., & Laborderie, A. (2005). The impact of landsat satellite monitoring on conservation biology. *Environ Monit Assess*, 106(1-3), 81-101.
- Louis, G. E., & Magpili, L. M. (2002). Representing inequities in the distribution of socio-economic benefits and environmental risk. *Environ Monit Assess*, 79(2), 101-119.
- Ma, M. G., Wang, X. M., & Cheng, G. D. (2003). Study on the oasis corridor landscape in the arid regions based on RS and GIS methods--application of Jinta Oasis, China. *J Environ Sci (China)*, 15(2), 193-198.
- MacLeod, M., Woodfine, D. G., Mackay, D., McKone, T., Bennett, D., & Maddalena, R. (2001). BETR North America: a regionally segmented multimedia contaminant fate model for North America. *Environ Sci Pollut Res Int*, 8(3), 156-163.
- Marsili-Libelli, S., Caporali, E., Arrighi, S., & Becattelli, C. (2001). A georeferenced river quality model. *Water Sci Technol*, 43(7), 223-230.
- Mertens, B., Forni, E., & Lambin, E. F. (2001). Prediction of the impact of logging activities on forest cover: a case-study in the east province of Cameroon. *J Environ Manage*, 62(1), 21-36.

- Miller, C., Chudek, P., & Babcock, S. (2003). A comparison of wellhead protection area delineation methods for public drinking water systems in Whatcom county, Washington. *J Environ Health*, 66(2), 17-26, 44.
- Miranda, M. L., & Dolinoy, D. C. (2005). Using GIS-based approaches to support research on neurotoxicants and other children's environmental health threats. *Neurotoxicology*, 26(2), 223-228.
- Molesworth, A. M., Harrison, R., Theakston, R. D., & Lalloo, D. G. (2003). Geographic Information System mapping of snakebite incidence in northern Ghana and Nigeria using environmental indicators: a preliminary study. *Trans R Soc Trop Med Hyg*, 97(2), 188-192.
- Nepal, S. K., & Nepal, S. A. (2004). Visitor impacts on trails in the Sagarmatha (Mt. Everest) National Park, Nepal. *Ambio*, 33(6), 334-340.
- Ostwald, M. (2002). GIS-based support tool system for decision-making regarding local forest protection: illustrations from Orissa, India. *Environ Manage*, 30(1), 35-45.
- Sandstrom, P., Pahlen, T. G., Edenius, L., Tommervik, H., Hagner, O., Hemberg, L., et al. (2003). Conflict resolution by participatory management: remote sensing and GIS as tools for communicating land-use needs for reindeer herding in northern Sweden. *Ambio*, 32(8), 557-567.
- Schiff, K. C., Weisberg, S. B., & Dorsey, J. H. (2001). Microbiological monitoring of marine recreational waters in southern California. *Environ Manage*, 27(1), 149-157.
- Shi, H., & Singh, A. (2003). Status and interconnections of selected environmental issues in the global coastal zones. *Ambio*, 32(2), 145-152.
- Svensson, M. K., Thorsson, S., & Lindqvist, S. (2003). A geographical information system model for creating bioclimatic maps - examples from a high, mid-latitude city. *Int J Biometeorol*, 47(2), 102-112.
- Turner, J., & Klaus, R. (2005). Coral reefs of the Mascarenes, Western Indian Ocean. *Philos Transact A Math Phys Eng Sci*, 363(1826), 229-250.
- van der Merwe, J. H., & Lohrentz, G. (2001). Demarcating coastal vegetation buffers with multicriteria evaluation and GIS at Saldanha Bay, South Africa. *Ambio*, 30(2), 89-95.
- Wang, X. (2001). Integrating water-quality management and land-use planning in a watershed context. *J Environ Manage*, 61(1), 25-36.
- Webb, E. L., & Thiha. (2002). Integrating social preference in GIS-aided planning for forestry and conservation activities: a case study from rural SE Asia. *Environ Manage*, 30(2), 183-198.

Pollution (air)

Andre, M., & Villanova, A. (2004). Characterisation of an urban bus network for environmental purposes. *Sci Total Environ*, 334-335, 85-99.

Bellander, T., Berglind, N., Gustavsson, P., Jonson, T., Nyberg, F., Pershagen, G., et al. (2001). Using geographic information systems to assess individual historical exposure to air pollution from traffic and house heating in Stockholm. *Environ Health Perspect*, 109(6), 633-639.

Boudet, C., Zmirou, D., & Vestri, V. (2001). Can one use ambient air concentration data to estimate personal and population exposures to particles? An approach within the European EXPOLIS study. *Sci Total Environ*, 267(1-3), 141-150.

Brauer, M., Hoek, G., van Vliet, P., Meliefste, K., Fischer, P., Gehring, U., et al. (2003). Estimating long-term average particulate air pollution concentrations: application of traffic indicators and geographic information systems. *Epidemiology*, 14(2), 228-239.

Briggs, D. (2005). The role of GIS: coping with space (and time) in air pollution exposure assessment. *J Toxicol Environ Health A*, 68(13-14), 1243-1261.

Brody, S. D., Peck, B. M., & Highfield, W. E. (2004). Examining localized patterns of air quality perception in Texas: a spatial and statistical analysis. *Risk Anal*, 24(6), 1561-1574.

Brunekreef, B. (2003). Design of cohort studies for air pollution health effects. *J Toxicol Environ Health A*, 66(16-19), 1723-1729.

Chakraborty, J. (2001). Acute exposure to extremely hazardous substances: an analysis of environmental equity. *Risk Anal*, 21(5), 883-895.

Cicero-Fernandez, P., Torres, V., Rosales, A., Cesar, H., Dorland, K., Munoz, R., et al. (2001). Evaluation of human exposure to ambient PM10 in the metropolitan area of Mexico City using a GIS-based methodology. *J Air Waste Manag Assoc*, 51(11), 1586-1593.

Crabbe, H., Barber, A., Bayford, R., Hamilton, R., Jarrett, D., & Machin, N. (2004). The use of a European telemedicine system to examine the effects of pollutants and allergens on asthmatic respiratory health. *Sci Total Environ*, 334-335, 417-426.

Dolinoy, D. C., & Miranda, M. L. (2004). GIS modeling of air toxics releases from TRI-reporting and non-TRI-reporting facilities: impacts for environmental justice. *Environ Health Perspect*, 112(17), 1717-1724.

Dubois, G., & De Cort, M. (2001). Mapping 137Cs deposition: data validation methods and data interpretation. *J Environ Radioact*, 53(3), 271-289.

- Gehring, U., Cyrus, J., Sedlmeir, G., Brunekreef, B., Bellander, T., Fischer, P., et al. (2002). Traffic-related air pollution and respiratory health during the first 2 yrs of life. *Eur Respir J*, 19(4), 690-698.
- Goswami, E., Larson, T., Lumley, T., & Liu, L. J. (2002). Spatial characteristics of fine particulate matter: identifying representative monitoring locations in Seattle, Washington. *J Air Waste Manag Assoc*, 52(3), 324-333.
- Gulliver, J., & Briggs, D. J. (2005). Time-space modeling of journey-time exposure to traffic-related air pollution using GIS. *Environ Res*, 97(1), 10-25.
- Heinrich, J., Gehring, U., Cyrus, J., Brauer, M., Hoek, G., Fischer, P., et al. (2005). Exposure to traffic related air pollutants: self reported traffic intensity versus GIS modelled exposure. *Occup Environ Med*, 62(8), 517-523.
- Hoek, G., Fischer, P., Van Den Brandt, P., Goldbohm, S., & Brunekreef, B. (2001). Estimation of long-term average exposure to outdoor air pollution for a cohort study on mortality. *J Expo Anal Environ Epidemiol*, 11(6), 459-469.
- Lange, J. L., Schwartz, D. A., Doebbeling, B. N., Heller, J. M., & Thorne, P. S. (2002). Exposures to the Kuwait oil fires and their association with asthma and bronchitis among gulf war veterans. *Environ Health Perspect*, 110(11), 1141-1146.
- LePoire, D. J., Arnish, J. J., Chen, S. Y., Faillace, E. R., Yuan, Y. C., & Schmidt, D. W. (2001). MILDOS uranium milling dose assessment code update. *Health Phys*, 81(5 Suppl), S78-84.
- Levy, J. I., Houseman, E. A., Spengler, J. D., Loh, P., & Ryan, L. (2001). Fine particulate matter and polycyclic aromatic hydrocarbon concentration patterns in Roxbury, Massachusetts: a community-based GIS analysis. *Environ Health Perspect*, 109(4), 341-347.
- Lindley, S. J., & Crabbe, H. (2004). What lies beneath?--issues in the representation of air quality management data for public consumption. *Sci Total Environ*, 334-335, 307-325.
- Pikhart, H., Bobak, M., Gorynski, P., Wojtyniak, B., Danova, J., Celko, M. A., et al. (2001). Outdoor sulphur dioxide and respiratory symptoms in Czech and Polish school children: a small-area study (SAVIAH). Small-Area Variation in Air Pollution and Health. *Int Arch Occup Environ Health*, 74(8), 574-578.
- Reungoat, P., Chiron, M., & Momas, I. (2004). [Assessment of exposure to traffic pollution in epidemiological studies: a review]. *Rev Epidemiol Sante Publique*, 52(3), 271-296.
- Ryan, P. H., LeMasters, G., Biagini, J., Bernstein, D., Grinshpun, S. A., Shukla, R., et al. (2005). Is it traffic type, volume, or distance? Wheezing in infants living near truck and bus traffic. *J Allergy Clin Immunol*, 116(2), 279-284.

Scoggins, A. (2004). Does air pollution pose a public health problem for New Zealand? *Aust N Z J Public Health*, 28(1), 16-19.

Scoggins, A., Kjellstrom, T., Fisher, G., Connor, J., & Gimson, N. (2004). Spatial analysis of annual air pollution exposure and mortality. *Sci Total Environ*, 321(1-3), 71-85.

Venn, A. J., Lewis, S. A., Cooper, M., Hubbard, R., & Britton, J. (2001). Living near a main road and the risk of wheezing illness in children. *Am J Respir Crit Care Med*, 164(12), 2177-2180.

Pollution (other)

Bertolo, A., & Verdi, L. (2001). Validation of a geographic information system for the evaluation of the soil radon exhalation potential in South-Tyrol and Veneto, Italy. *Radiat Prot Dosimetry*, 97(4), 321-324.

Brody, J. G., Vorhees, D. J., Melly, S. J., Swedis, S. R., Drivas, P. J., & Rudel, R. A. (2002). Using GIS and historical records to reconstruct residential exposure to large-scale pesticide application. *J Expo Anal Environ Epidemiol*, 12(1), 64-80.

Franco-Marina, F., Villalba-Caloca, J., Segovia, N., & Tavera, L. (2003). Spatial indoor radon distribution in Mexico City. *Sci Total Environ*, 317(1-3), 91-103.

Hsu, E. B., Grabowski, J. G., Chotani, R. A., Winslow, J. A., Alves, D. W., & VanRooyen, M. J. (2002). Effects on local emergency departments of large-scale urban chemical fire with hazardous materials spill. *Prehospital Disaster Med*, 17(4), 196-201.

Ielsch, G., Ferry, C., Tymen, G., & Robe, M. C. (2002). Study of a predictive methodology for quantification and mapping of the radon-222 exhalation rate. *J Environ Radioact*, 63(1), 15-33.

Kaminska, I. A., Oldak, A., & Turski, W. A. (2004). Geographical Information System (GIS) as a tool for monitoring and analysing pesticide pollution and its impact on public health. *Ann Agric Environ Med*, 11(2), 181-184.

Kontos, T. D., Komilis, D. P., & Halvadakis, C. R. (2003). Siting MSW landfills on Lesbos island with a GIS-based methodology. *Waste Manag Res*, 21(3), 262-277.

Li, J., & Hao, J. (2003). Application of intake fraction to population exposure estimates in Hunan Province of China. *J Environ Sci Health A Tox Hazard Subst Environ Eng*, 38(6), 1041-1054.

Maantay, J. (2002). Mapping environmental injustices: pitfalls and potential of geographic information systems in assessing environmental health and equity. *Environ Health Perspect*, 110 Suppl 2, 161-171.

- Macey, G. P., Her, X., Reibling, E. T., & Ericson, J. (2001). An investigation of environmental racism claims: testing environmental management approaches with a geographic information system. *Environ Manage*, 27(6), 893-907.
- Miranda, M. L., Dolinoy, D. C., & Overstreet, M. A. (2002). Mapping for prevention: GIS models for directing childhood lead poisoning prevention programs. *Environ Health Perspect*, 110(9), 947-953.
- Petropoulos, N. P., Anagnostakis, M. J., Hinis, E. P., & Simopoulos, S. E. (2001). Geographical mapping and associated fractal analysis of the long-lived Chernobyl fallout radionuclides in Greece. *J Environ Radioact*, 53(1), 59-66.
- Reissman, D. B., Staley, F., Curtis, G. B., & Kaufmann, R. B. (2001). Use of geographic information system technology to aid Health Department decision making about childhood lead poisoning prevention activities. *Environ Health Perspect*, 109(1), 89-94.
- Roberts, J. R., Hulsey, T. C., Curtis, G. B., & Reigart, J. R. (2003). Using geographic information systems to assess risk for elevated blood lead levels in children. *Public Health Rep*, 118(3), 221-229.
- Rojas, M., Espinosa, C., & Seijas, D. (2003). [Association between blood lead and sociodemographic parameters among children]. *Rev Saude Publica*, 37(4), 503-509.
- Royster, M. O., Hilborn, E. D., Barr, D., Carty, C. L., Rhoney, S., & Walsh, D. (2002). A pilot study of global positioning system/geographical information system measurement of residential proximity to agricultural fields and urinary organophosphate metabolite concentrations in toddlers. *J Expo Anal Environ Epidemiol*, 12(6), 433-440.
- Rybach, L., Bachler, D., Bucher, B., & Schwarz, G. (2002). Radiation doses of Swiss population from external sources. *J Environ Radioact*, 62(3), 277-286.
- Stellman, J. M., Stellman, S. D., Weber, T., Tomasallo, C., Stellman, A. B., & Christian, R., Jr. (2003). A geographic information system for characterizing exposure to Agent Orange and other herbicides in Vietnam. *Environ Health Perspect*, 111(3), 321-328.
- Stellman, S. D., & Stellman, J. M. (2004). Exposure opportunity models for Agent Orange, dioxin, and other military herbicides used in Vietnam, 1961-1971. *J Expo Anal Environ Epidemiol*, 14(4), 354-362.
- Sudakin, D. L., Horowitz, Z., & Giffin, S. (2002). Regional variation in the incidence of symptomatic pesticide exposures: applications of geographic information systems. *J Toxicol Clin Toxicol*, 40(6), 767-773.
- Zh, H. C., Charlet, J. M., & Poffijn, A. (2001). Radon risk mapping in southern Belgium: an application of geostatistical and GIS techniques. *Sci Total Environ*, 272(1-3), 203-210.

Pollution (water)

Cifuentes, E., Suarez, L., Solano, M., & Santos, R. (2002). Diarrheal diseases in children from a water reclamation site in Mexico city. *Environ Health Perspect*, 110(10), A619-624.

Dangendorf, F., Herbst, S., Reintjes, R., & Kistemann, T. (2002). Spatial patterns of diarrhoeal illnesses with regard to water supply structures--a GIS analysis. *Int J Hyg Environ Health*, 205(3), 183-191.

Hales, S., Black, W., Skelly, C., Salmond, C., & Weinstein, P. (2003). Social deprivation and the public health risks of community drinking water supplies in New Zealand. *J Epidemiol Community Health*, 57(8), 581-583.

Hudak, P. F., & Sanmanee, S. (2003). Spatial patterns of nitrate, chloride, sulfate, and fluoride concentrations in the Woodbine Aquifer of north-central Texas. *Environ Monit Assess*, 82(3), 311-320.

Kistemann, T., Dangendorf, F., & Exner, M. (2001). A geographical information system (GIS) as a tool for microbial risk assessment in catchment areas of drinking water reservoirs. *Int J Hyg Environ Health*, 203(3), 225-233.

Reif, J. S., Burch, J. B., Nuckols, J. R., Metzger, L., Ellington, D., & Anger, W. K. (2003). Neurobehavioral effects of exposure to trichloroethylene through a municipal water supply. *Environ Res*, 93(3), 248-258.

Schowaneck, D., & Webb, S. (2002). Exposure simulation for pharmaceuticals in European surface waters with GREAT-ER. *Toxicol Lett*, 131(1-2), 39-50.

Segala, C., Poizeau, D., & Mace, J. M. (2003). [Odors and health: a descriptive epidemiological study around a wastewater treatment plant]. *Rev Epidemiol Sante Publique*, 51(2), 201-214.

Swartz, C. H., Rudel, R. A., Kachajian, J. R., & Brody, J. G. (2003). Historical reconstruction of wastewater and land use impacts to groundwater used for public drinking water: exposure assessment using chemical data and GIS. *J Expo Anal Environ Epidemiol*, 13(5), 403-416.

Thunqvist, E. L. (2004). Regional increase of mean chloride concentration in water due to the application of deicing salt. *Sci Total Environ*, 325(1-3), 29-37.

Economics/Urban

Public Policy and Politics

Benini, A. A., Conley, C. E., Shdeed, R., Spurway, K., & Yarmoshuk, M. (2003). Integration of different data bodies for humanitarian decision support: an example from mine action. *Disasters*, 27(4), 288-304.

Cowper, D., Yu, W., Kuebeler, M., Kubal, J. D., Manheim, L. M., & Ripley, B. A. (2004). Using GIS in government: an overview of the VHA's Healthcare Atlas, FY-2000. *J Med Syst*, 28(3), 257-269.

Transportation

Guzzetti, F., Reichenbach, P., & Ghigi, S. (2004). Rockfall hazard and risk assessment along a transportation corridor in the Nera Valley, central Italy. *Environ Manage*, 34(2), 191-208.

Urban Planning and Community

Buckeridge, D. L., Mason, R., Robertson, A., Frank, J., Glazier, R., Purdon, L., et al. (2002). Making health data maps: a case study of a community/university research collaboration. *Soc Sci Med*, 55(7), 1189-1206.

English, P. B., Kharrazi, M., Davies, S., Scalf, R., Waller, L., & Neutra, R. (2003). Changes in the spatial pattern of low birth weight in a southern California county: the role of individual and neighborhood level factors. *Soc Sci Med*, 56(10), 2073-2088.

Faruque, F. S., Lofton, S. P., Doddato, T. M., & Mangum, C. (2003). Utilizing Geographic Information Systems in community assessment and nursing research. *J Community Health Nurs*, 20(3), 179-191.

Green, R. S., Smorodinsky, S., Kim, J. J., McLaughlin, R., & Ostro, B. (2004). Proximity of California public schools to busy roads. *Environ Health Perspect*, 112(1), 61-66.

Hong, S. K., Song, I. J., Kim, H. O., & Lee, E. K. (2003). Landscape pattern and its effect on ecosystem functions in Seoul Metropolitan area: urban ecology on distribution of the naturalized plant species. *J Environ Sci (China)*, 15(2), 199-204.

Le Lay, G., Clergeau, P., & Hubert-Moy, L. (2001). Computerized map of risk to manage wildlife species in urban areas. *Environ Manage*, 27(3), 451-461.

Novotny, V., Clark, D., Griffin, R. J., & Booth, D. (2001). Risk based urban watershed management under conflicting objectives. *Water Sci Technol*, 43(5), 69-78.

Schneider, R. J., Ryznar, R. M., & Khattak, A. J. (2004). An accident waiting to happen: a spatial approach to proactive pedestrian planning. *Accid Anal Prev*, 36(2), 193-211.

Thums, C., & Farago, M. (2001). Investigating urban geochemistry using Geographical Information Systems. *Sci Prog*, 84(Pt 3), 183-204.

Whitman, S., Silva, A., Shah, A., & Ansell, D. (2004). Diversity and disparity: GIS and small-area analysis in six Chicago neighborhoods. *J Med Syst*, 28(4), 397-411.

Zhao, B., Nakagoshi, N., Chen, J. K., & Kong, L. Y. (2003). The impact of urban planning on land use and land cover in Pudong of Shanghai, China. *J Environ Sci (China)*, 15(2), 205-214.

Social Science

Anthropology

Chaplin, G. (2004). Geographic distribution of environmental factors influencing human skin coloration. *Am J Phys Anthropol*, 125(3), 292-302.

Pavesi, A. (2004). Detecting Traces of Prehistoric Human Migrations by Geographic Synthetic Maps of Polyomavirus JC. *J Mol Evol*, 58(3), 304-313.

Ungar, P. (2004). Dental topography and diets of *Australopithecus afarensis* and early Homo. *J Hum Evol*, 46(5), 605-622.

Crime

Santos, S. M., Barcellos, C., Carvalho, M. S., & Flores, R. (2001). [Spatial clusters detection of violent deaths in Porto Alegre, Rio Grande do Sul, Brazil, 1996]. *Cad Saude Publica*, 17(5), 1141-1151.

Deprivation

Niggebrugge, A., Haynes, R., Jones, A., Lovett, A., & Harvey, I. (2005). The index of multiple deprivation 2000 access domain: a useful indicator for public health? *Soc Sci Med*, 60(12), 2743-2753.

Methods/Applications

Agent Based Modeling

Evans, T. P., & Kelley, H. (2004). Multi-scale analysis of a household level agent-based model of landcover change. *J Environ Manage*, 72(1-2), 57-72.

Atlases

Benach, J., Yasui, Y., Borrell, C., Rosa, E., Pasarin, M. I., Benach, N., et al. (2003). Examining geographic patterns of mortality: the atlas of mortality in small areas in Spain (1987-1995). *Eur J Public Health*, 13(2), 115-123.

Geocoding

Bonner, M. R., Han, D., Nie, J., Rogerson, P., Vena, J. E., & Freudenheim, J. L. (2003). Positional accuracy of geocoded addresses in epidemiologic research. *Epidemiology*, 14(4), 408-412.

Hurley, S. E., Saunders, T. M., Nivas, R., Hertz, A., & Reynolds, P. (2003). Post office box addresses: a challenge for geographic information system-based studies. *Epidemiology*, 14(4), 386-391.

McElroy, J. A., Remington, P. L., Trentham-Dietz, A., Robert, S. A., & Newcomb, P. A. (2003). Geocoding addresses from a large population-based study: lessons learned. *Epidemiology*, 14(4), 399-407.

Rushton, G., Peleg, I., Banerjee, A., Smith, G., & West, M. (2004). Analyzing geographic patterns of disease incidence: rates of late-stage colorectal cancer in Iowa. *J Med Syst*, 28(3), 223-236.

Skaba, D. A., Carvalho, M. S., Barcellos, C., Martins, P. C., & Terron, S. L. (2004). [Geoprocessing of health data: treatment of information on addresses]. *Cad Saude Publica*, 20(6), 1753-1756.

Whitsel, E. A., Rose, K. M., Wood, J. L., Henley, A. C., Liao, D., & Heiss, G. (2004). Accuracy and repeatability of commercial geocoding. *Am J Epidemiol*, 160(10), 1023-1029.

Point Pattern

Cockings, S., Dunn, C. E., Bhopal, R. S., & Walker, D. R. (2004). Users' perspectives on epidemiological, GIS and point pattern approaches to analysing environment and health data. *Health Place*, 10(2), 169-182.

Spatial Analysis/Statistics Issues

Banerjee, S., & Dey, D. K. (2005). Semiparametric proportional odds models for spatially correlated survival data. *Lifetime Data Anal*, 11(2), 175-191.

Fang, Z., Kulldorff, M., & Gregorio, D. I. (2004). Brain cancer mortality in the United States, 1986 to 1995: a geographic analysis. *Neuro-oncol*, 6(3), 179-187.

Law, D. C., Serre, M. L., Christakos, G., Leone, P. A., & Miller, W. C. (2004). Spatial analysis and mapping of sexually transmitted diseases to optimise intervention and prevention strategies. *Sex Transm Infect*, 80(4), 294-299.

Luke, D. A. (2005). Getting the big picture in community science: methods that capture context. *Am J Community Psychol*, 35(3-4), 185-200.

Slachtova, H., Machova, T., Tomaskova, H., & Michalik, J. (2003). Evaluation of the response of a questionnaire study by using the GIS and standard statistical methods. *Cent Eur J Public Health*, 11(2), 79-86.

Technology

Appelrath, H. J., Friebe, J., Grawunder, M., & Wellmann, I. (2001). The need for open geographical information systems in medicine. *Methods Inf Med*, 40(2), 69-73.

Croner, C. M. (2003). Public health, GIS, and the internet. *Annu Rev Public Health*, 24, 57-82.

da Silva, N. N., da Cunha, T. N., & Quintanilha, J. A. (2003). [Master sample and geoprocessing: technologies for household surveys]. *Rev Saude Publica*, 37(4), 494-502.

Elgethun, K., Fenske, R. A., Yost, M. G., & Palcisko, G. J. (2003). Time-location analysis for exposure assessment studies of children using a novel global positioning system instrument. *Environ Health Perspect*, 111(1), 115-122.

Loomis, J. M., Klatzky, R. L., & Golledge, R. G. (2001). Navigating without vision: basic and applied research. *Optom Vis Sci*, 78(5), 282-289.

Rushton, G. (2003). Public health, GIS, and spatial analytic tools. *Annu Rev Public Health*, 24, 43-56.

Witte, T. H., & Wilson, A. M. (2005). Accuracy of WAAS-enabled GPS for the determination of position and speed over ground. *J Biomech*, 38(8), 1717-1722.

Zhou, X. N., Hu, X. S., Yang, G. J., Sun, N. S., Wang, T. P., Malone, J., et al. (2003). [Establishment of minimum medical geographic information systems database in China]. *Zhonghua Liu Xing Bing Xue Za Zhi*, 24(4), 253-256.

WebGIS

Paolino, L., Sebillo, M., & Cringoli, G. (2005). Geographical Information Systems and on-line GIServices for health data sharing and management. *Parassitologia*, 47(1), 171-175.